

ON THE ASYMPTOTIC DISTRIBUTION OF ZEROS OF MODULAR FORMS

ZE'EV RUDNICK

ABSTRACT. We study the distribution of zeros of holomorphic modular forms. Assuming the Generalized Riemann Hypothesis we show that the zeros of Hecke eigenforms for the modular group become equidistributed with respect to the hyperbolic measure on the modular domain as the weight grows.

1. INTRODUCTION

1.1. Our purpose in this note is to study the limiting distribution of zeros of modular forms. We review some definitions: A modular form of weight k for $SL_2(\mathbb{Z})$ is a holomorphic function on the upper half-plane $\mathbb{H}$, transforming as $f(\frac{az+b}{cz+d}) = (cz+d)^k f(z)$, for all $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{Z})$ (this forces k to be even), and “holomorphic at the cusp” (see § 2.1). A form is *cuspidal* if it vanishes at the cusp. For a modular form of weight k , let $\nu(f)$ be the number of inequivalent zeros of f in $\mathbb{H}$, with the convention that a zero at z is counted with weight $w(z)$ inverse to the number of elements of $SL_2(\mathbb{Z})/\{\pm I\}$ fixing z . Then $\nu(f) \leq k/12$.

For a sequence of modular forms, where we assume that the number of inequivalent zeros tends to infinity, we would like to examine the manner in which the resulting configuration of zeros is distributed in the modular domain $SL_2(\mathbb{Z}) \backslash \mathbb{H}$.

1.2. **Example: Eisenstein series.** The Eisenstein series are non-cuspidal modular forms of weight $k > 2$, given by the sum

$$E_k(z) = \frac{1}{2} \sum_{(c,d)=1} \frac{1}{(cz+d)^k}.$$

F.K.C. Rankin and Swinnerton-Dyer [13] showed that all zeros (in the fundamental domain) of E_k lie on boundary arc $\{|z| = 1\}$. Moreover, as

Date: June 14, 2005.

Partially supported by grant number 2002088 from the United States-Israel Binational Science Foundation (BSF), Jerusalem, Israel.

$k \rightarrow \infty$, the zeros become uniformly distributed on the unit arc. R.A. Rankin [14] gave a similar result for the (cuspidal) Poincaré series

$$G_k(z, m) = \sum_{\gamma \in \Gamma_\infty \backslash \Gamma} \frac{\exp(2\pi i m \gamma z)}{(cz + d)^k}$$

with $m \geq 1$ fixed, $k \gg m$. See [1] for yet another such example.

1.3. Example: higher Weierstrass points. An example in a slightly different setting are the higher Weierstrass points on a (compact) Riemann surface C . Let $S_k(C)$ be the space of holomorphic forms of weight k on C , $d := \dim S_k(C)$, and let $f_1, \dots, f_d$ be a basis for $S_k(C)$. The Wronskian is defined as

$$\text{Wr}_k(\{f_j\}; z) := \det(f_j^{(i)})_{1 \leq i, j \leq d}$$

It is a modular form of weight $d(k + d - 1)$.

A k -th order Weierstrass point is a zero of Wr_k . Equivalently, these are points of C where there is a nonzero form which vanishes to order $\geq \dim S_k(C)$.

It was conjectured by Bers, and proved by Olsen [10] that as $k \rightarrow \infty$, these become dense in C . Mumford (see page 11 of [7]) and Neeman [8] showed that in fact these become equidistributed with respect to the Arakelov (or Bergmann) measure on C , which arises from the metric on C gotten by embedding the curve in its Jacobian and pulling back the flat metric.

1.4. In contrast to these examples, we consider the case of cuspidal Hecke eigenforms. Recall that the Hecke operators

$$T_n f(z) := \frac{1}{n} \sum_{ad=n} a^k \sum_{b \pmod d} f\left(\frac{az + b}{d}\right)$$

act on the space of cusp forms of weight k , commute with each other and are self-adjoint with respect to the Petersson inner product

$$\langle f, g \rangle := \int_{SL_2(\mathbb{Z}) \backslash \mathbb{H}} f(z) \overline{g(z)} y^k \frac{dx dy}{y^2}$$

Thus the space of cusp forms admits a basis consisting of joint eigenfunctions of all Hecke operators (the Eisenstein series E_k is also an eigenfunction). The cuspidal Hecke eigenfunctions have a simple zero at the cusp. For these Hecke eigenforms, we have:

Theorem 1. *Assume the Generalized Riemann Hypothesis (GRH). Let $\{f_k\}$ be a sequence of cuspidal Hecke eigenforms, then as $k \rightarrow \infty$ their*

zeros are equidistributed in $SL_2(\mathbb{Z}) \backslash \mathbb{H}$ with respect to the normalized hyperbolic measure $dV(z) = \frac{3}{\pi} \frac{dx dy}{y^2}$.

Equidistribution here means that for any nice compact subset $\Omega \subset SL_2(\mathbb{Z}) \backslash \mathbb{H}$, the proportion of (inequivalent) zeros of such a Hecke eigenform which lie in Ω is asymptotically the relative area of Ω :

$$\frac{1}{\nu(f)} \sum_{z \in \Omega} w(z) \text{ord}_z(f) \sim \int_{\Omega} dV(z)$$

1.5. Theorem 1 is the outgrowth of some ideas from the theory of “quantum chaos”. The mechanism is that equidistribution of the zeros of a sequence of forms f_k is implied by equidistribution of the “masses” $y^k |f_k(z)|^2 dV(z)$ of the forms. This idea was discovered by Nonnenmacher and Voros [9] in the context of quantum maps. Around the same time a very general result of this sort was obtained by Shiffman and Zelditch [18] for zeros of sections of high powers of a positive holomorphic hermitian line bundle over any compact complex manifold ([9] deal with curves of genus one). All these are in a compact setting. The analogous result in our case is

Theorem 2. *Suppose that $\{f_k\}$ is a sequence of L^2 -normalized cusp forms (f_k of weight k) for which the bulk of zeros lie in the fundamental domain: $\nu(f_k) \sim k/12$. Assume that for some $c > 0$, we have*

$$(1) \quad y^k |f_k(z)|^2 dV(z) \xrightarrow{w} c \cdot dV(z)$$

Then the zeros of f_k are equidistributed with respect to $dV(z)$.

Here $\xrightarrow{w}$ denotes weak convergence when we test against compactly supported functions.

The proof of Theorem 2, given in Section 3, follows closely the argument of [9, 18], with care taken to handle the complications due to the lack of compactness of the modular domain. We require the hypothesis (1), which is weaker than equidistribution of the masses as it allows “leakage” of some of the mass at the cusp (which cannot happen in a compact setting). This slightly weaker version of equidistribution of masses is what Lindenstrauss [5] proved unconditionally for the analogous case of Maass forms, though he cannot exclude $c = 0$. In the holomorphic case we do not have an unconditional proof of (1). However it has been known for some time to follow (with $c = 1$) from GRH [16, 21]. This is sketched in section 4.

Acknowledgements: This manuscript is based on notes written in October 1999, while enjoying the hospitality of the Institute of Advanced Studies in Princeton. I have benefited from several discussions with Peter Sarnak, Misha Sodin and Steve Zelditch.

2. POTENTIAL THEORY ON $\Gamma \backslash \mathbb{H}$

2.1. Preliminaries. We review some definitions: A modular form of weight k for $SL_2(\mathbb{Z})$ is a holomorphic function on the upper half-plane $\mathbb{H}$, transforming as

$$f\left(\frac{az+b}{cz+d}\right) = (cz+d)^k f(z), \quad \text{for all } \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{Z})$$

(this forces k to be even), and *holomorphic at the cusp*. This means the following: Since $f(z)$ is periodic, it can be expressed as a holomorphic function $\tilde{f}(q)$ of $q = e^{2\pi iz}$ in the punctured disk $0 < |q| < 1$. The requirement to be holomorphic at the cusp means that it extends to a holomorphic function at $q = 0$. A form is *cuspidal* if it vanishes at the cusp, that is $\tilde{f}(0) = 0$. The order of vanishing at the cusp $\text{ord}_\infty(f)$ is defined as the order of vanishing at $q = 0$ of $\tilde{f}(q)$.

We denote by $\Gamma = SL_2(\mathbb{Z})/\{\pm I\}$ and speak interchangeably about modular forms for Γ and for $SL_2(\mathbb{Z})$. For each $z \in \mathbb{H}$ let $\Gamma_z = \{\gamma \in \Gamma : \gamma z = z\}$ be the stabilizer in Γ of z , and set

$$w(z) := \frac{1}{\#\Gamma_z} = \begin{cases} 1/2, & z \in \Gamma i \\ 1/3, & z \in \Gamma e^{2\pi i/3} \\ 1, & \text{otherwise} \end{cases}$$

For a modular form of weight k , let

$$\nu(f) = \sum_{z \in SL_2(\mathbb{Z}) \backslash \mathbb{H}} w(z) \text{ord}_z(f)$$

be the (weighted) number of Γ -inequivalent zeros of f in $\mathbb{H}$. Then $\nu(f) \leq \frac{k}{12}$, in fact $\nu(f) + \text{ord}_\infty(f) = \frac{k}{12}$. Note that $\nu(f) = 0$ only for powers of the modular discriminant.

2.2. For a smooth compactly supported function $\phi \in C_c^\infty(\mathbb{H})$ on the upper half-plane $\mathbb{H}$ set

$$F_\phi(z) := \sum_{\gamma \in \Gamma} \phi(\gamma z) \in C_c^\infty(\Gamma \backslash \mathbb{H})$$

Note that

$$\int_{\Gamma \backslash \mathbb{H}} F_\phi(z) \frac{dx dy}{y^2} = \int_{\mathbb{H}} \phi(z) dx dy$$

Let $dV(z) = \frac{1}{\text{vol}(\Gamma \backslash \mathbb{H})} \frac{dx dy}{y^2}$ be the normalized hyperbolic measure on the quotient $\Gamma \backslash \mathbb{H}$.

Lemma 2.1. *Let f be a (weakly holomorphic) modular form of weight k for Γ and let $\{z_j\}$ be a set of Γ -inequivalent zeros of f in $\mathbb{H}$. Then*

$$(2) \quad \sum_j w(z_j) F_\phi(z_j) = k \frac{\text{vol}(\Gamma \backslash \mathbb{H})}{4\pi} \int_{\Gamma \backslash \mathbb{H}} F_\phi(z) dV(z) \\ + \frac{1}{2\pi} \int_{\mathbb{H}} \log(y^{k/2} |f(z)|) \Delta \phi(z) dx dy$$

Proof. Let $S = \{z \in \mathbb{H} : f(z) = 0\}$ be the set of zeros of the form f . It is a discrete set of points, which is stable under Γ ($\gamma S = S$ for all $\gamma \in \Gamma$). For any such S we have

$$(3) \quad \sum_{z \in \Gamma \backslash S} w(z) F_\phi(z) = \sum_{s \in S} \phi(s)$$

Indeed,

$$\begin{aligned} \sum_{s \in S} \phi(s) &= \sum_{z \in \Gamma \backslash S} \sum_{s \in \Gamma z} \phi(s) \\ &= \sum_{z \in \Gamma \backslash S} \sum_{\gamma \in \Gamma / \Gamma_z} \phi(\gamma z) \\ &= \sum_{z \in \Gamma \backslash S} \frac{1}{\#\Gamma_z} \sum_{\gamma \in \Gamma} \phi(\gamma z) \\ &= \sum_{z \in \Gamma \backslash S} \frac{1}{\#\Gamma_z} F_\phi(z) \end{aligned}$$

as required.

We recall that the electrostatic potential for a point charge in the plane is $\frac{1}{2\pi} \log |z|$, that is for $\phi \in C_c^\infty(\mathbb{C})$ we have

$$\int_{\mathbb{C}} \frac{1}{2\pi} \log |z| \Delta \phi(z) dx dy = \phi(0)$$

Consequently, for the holomorphic function $f(z)$ on $\mathbb{H}$ we have

$$(4) \quad \int_{\mathbb{H}} \frac{1}{2\pi} \log |f(z)| \Delta \phi(z) dx dy = \sum_{s: f(s)=0} \phi(s)$$

for all $\phi \in C_c^\infty(\mathbb{H})$ (with multiple zeros repeated). Thus

$$\begin{aligned} \sum_j w(z_j) F_\phi(z_j) &= \frac{1}{2\pi} \int_{\mathbb{H}} \log |f(z)| \Delta \phi(z) dx dy \\ &= -\frac{1}{2\pi} \int_{\mathbb{H}} \log y^{k/2} \Delta \phi(z) dx dy + \frac{1}{2\pi} \int_{\mathbb{H}} \log(y^{k/2} |f(z)|) \Delta \phi(z) dx dy \end{aligned}$$

The first term above is transformed via integration by parts into

$$-\frac{1}{2\pi} \int_{\mathbb{H}} \Delta(\log y^{k/2}) \phi(z) dx dy = \frac{k}{4\pi} \int_{\mathbb{H}} \phi(z) \frac{dx dy}{y^2}$$

Note that

$$\int_{\mathbb{H}} \phi(z) \frac{dx dy}{y^2} = \int_{\Gamma \backslash \mathbb{H}} F_{\phi}(z) \frac{dx dy}{y^2} = \text{vol}(\Gamma \backslash \mathbb{H}) \int_{\Gamma \backslash \mathbb{H}} F_{\phi}(z) dV(z)$$

so that we get

$$-\frac{1}{2\pi} \int_{\mathbb{H}} \Delta(\log y^{k/2}) \phi(z) dx dy = k \frac{\text{vol}(\Gamma \backslash \mathbb{H})}{4\pi} \int_{\Gamma \backslash \mathbb{H}} F_{\phi}(z) dV(z)$$

as required. $\square$

Remark: We may reformulate this in Γ -invariant form as follows: Instead of the Euclidean Laplacian $\Delta = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}$ we use the hyperbolic Laplacian

$$\mathcal{L} = y^2 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right)$$

which is Γ -invariant, and instead of $|f(z)|$ we use $y^k |f(z)|^2$ which is Γ -invariant. Then

$$(5) \quad \sum_j w(z_j) F_{\phi}(z_j) = k \frac{\text{vol}(\Gamma \backslash \mathbb{H})}{4\pi} \int_{\Gamma \backslash \mathbb{H}} F_{\phi}(z) dV(z) \\ + \frac{\text{vol}(\Gamma \backslash \mathbb{H})}{2\pi} \int_{\Gamma \backslash \mathbb{H}} \log(y^{k/2} |f(z)|) \mathcal{L} F_{\phi}(z) \frac{dx dy}{y^2}$$

To derive (5) from Lemma 2.1, we transform the second term in (2) by noting that since $\frac{dx dy}{y^2}$ is Γ -invariant, we have

$$\begin{aligned} \frac{1}{2\pi} \int_{\mathbb{H}} \log(y^{k/2} |f(z)|) \mathcal{L} \phi(z) \frac{dx dy}{y^2} &= \frac{1}{2\pi} \int_{\Gamma \backslash \mathbb{H}} \log(y^{k/2} |f(z)|) \sum_{\gamma \in \Gamma} (\mathcal{L} \phi)(\gamma z) \frac{dx dy}{y^2} \\ &= \frac{1}{2\pi} \int_{\Gamma \backslash \mathbb{H}} \log(y^{k/2} |f(z)|) F_{\mathcal{L} \phi}(z) \frac{dx dy}{y^2} \\ &= \frac{\text{vol}(\Gamma \backslash \mathbb{H})}{2\pi} \int_{\Gamma \backslash \mathbb{H}} \log(y^{k/2} |f(z)|) F_{\mathcal{L} \phi}(z) dV(z) \end{aligned}$$

This proves (5) once we note that $F_{\mathcal{L} \phi} = \mathcal{L} F_{\phi}$ since $\mathcal{L}$ is Γ -invariant.

2.3. Background on sub-harmonic functions. Let $\Omega \subset \mathbb{C} = \mathbb{R}^2$ be an open connected set. An upper semi-continuous¹ function $u : \Omega \rightarrow [-\infty, \infty)$ is *sub-harmonic* if u is not identically $-\infty$ and for all $z \in \Omega$,

$$u(z) \leq \frac{1}{2\pi} \int_0^{2\pi} u(z + re^{i\theta}) d\theta$$

whenever the disc $\{w : |z - w| \leq r\}$ is contained in Ω .

Note: u is sub-harmonic $\Leftrightarrow \Delta u \geq 0$ as a distribution.

A fundamental example of a sub-harmonic function is $\log |f(z)|$ where $f(z)$ is holomorphic in Ω . Moreover we have

$$\Delta \log |f(z)| = 2\pi \sum_{f(z_j)=0} \delta(z - z_j)$$

We denote the space of subharmonic functions on Ω by $\mathcal{SH}(\Omega)$.

A basic compactness property of sub-harmonic functions is (c.f. [2], Thm 4.1.9):

Lemma 2.2 (Compactness property). *If $\{u_j\} \subset \mathcal{SH}(\Omega)$ are locally uniformly upper bounded (that is for all compact $K \subset \Omega$ there is c_K so that $u_j(z) \leq c_K$ for all $z \in K$ and all j) then either*

- (1) $u_j \rightarrow -\infty$ uniformly on compacta, OR
- (2) There is a subsequence u_{j_k} which converges weakly to some $u \in \mathcal{SH}(\Omega)$.
- (3) Moreover, in this case $\limsup u_j \leq u$ and $\limsup u_j = u$ almost everywhere.

The following is known as ‘‘Hartogs’ Lemma’’ ([15] Thm 3.4.3):

Lemma 2.3. *If $\{u_j\} \subset \mathcal{SH}(\Omega)$ are locally uniformly bounded above, and there is a continuous function $\phi : \Omega \rightarrow \mathbb{R}$ such that $\limsup u_j \leq \phi$, then $\max(u_j, \phi) \rightarrow \phi$ locally uniformly on Ω as $j \rightarrow \infty$.*

3. ERGODICITY OF EIGENFUNCTIONS IMPLIES EQUIDISTRIBUTION OF ZEROS

3.1. Proof of Theorem 2. Assume that there is some $c > 0$ so that $y^k |f_k|^2 dV \xrightarrow{w} cdV$ for a sequence of $k \rightarrow \infty$. We will show that necessarily the zeros $\{z_j\}$ of f_k become equidistributed relative to dV as $k \rightarrow \infty$. We need to show that for all $F \in C_c^\infty(\Gamma \backslash \mathbb{H})$ we have

$$\frac{1}{\nu(f_k)} \sum_j w(z_j) F(z_j) \sim \int_{\Gamma \backslash \mathbb{H}} F(z) dV(z), \quad k \rightarrow \infty$$

¹ $u : \Omega \rightarrow [-\infty, \infty)$ is upper semi-continuous if for all $c \in \mathbb{R}$, the set $\{z : u(z) < c\}$ is open.

It suffices to show this for F of the form $F(z) = F_\phi(z) = \sum_{\gamma \in \Gamma} \phi(\gamma z)$, $\phi \in C_c^\infty(\mathbb{H})$, since these are dense in $C_c^\infty(\Gamma \backslash \mathbb{H})$ with respect to the uniform topology.² For these we have the fundamental identity of Lemma 2.1

$$(6) \quad \frac{1}{\nu(f)} \sum_j w(z_j) F_\phi(z_j) = \frac{k/12}{\nu(f_k)} \int_{\Gamma \backslash \mathbb{H}} F_\phi(z) dV(z) + \mathcal{E}$$

where

$$\mathcal{E} = \frac{1}{2\pi\nu(f_k)} \int_{\mathbb{H}} \log(y^k |f_k(z)|^2) \Delta\phi(z) dx dy$$

Since we assume that $\nu(f_k) \sim k/12$, the main term is the desired one (the mean value of F_ϕ) and we need to show that $\mathcal{E} \rightarrow 0$ as $k \rightarrow \infty$, i. e. that for all $\phi \in C_c^\infty(\mathbb{H})$,

$$\int_{\mathbb{H}} \frac{1}{k} \log(y^k |f_k(z)|^2) \Delta\phi(z) dx dy \rightarrow 0$$

or that

$$(7) \quad \int_{\mathbb{H}} \frac{1}{k} \log(|f_k(z)|^2) \Delta\phi(z) dx dy \rightarrow \int_{\mathbb{H}} -\log y \Delta\phi(z) dx dy$$

Assume that this is false, that is there is some test function $\phi_0 \in C_c^\infty(\mathbb{H}/\Gamma)$ and a subsequence of the f_k (which we will continue to call f_k for notational ease) for which (7) fails. We will proceed to derive a contradiction.

We begin by listing some properties of $v_k := \frac{1}{k} \log |f_k^2|$. These are subharmonic in $\mathbb{H}/\Gamma$. By Proposition 3, we have $y^k |f_k(z)|^2 \ll k$ uniformly on compacta and so

$$(8) \quad v_k \leq -\log y + \frac{\log k + O(1)}{k}$$

thus

$$(9) \quad \limsup v_k \leq -\log y$$

locally uniformly. Thus the family $\{v_k\} \subset \mathcal{SH}$ is locally bounded above. By the compactness property (Lemma 2.2), this gives us two possibilities:

- i) $v_k \rightarrow -\infty$ locally uniformly, OR
- ii) $\{v_k\}$ has a weakly convergent subsequence.

We will dispose of both possibilities.

²To see this it suffices to show that for fixed ball $B \subset \mathbb{H}$, the space $\{F_\phi : \text{supp } \phi \subset B\}$ is dense in $C(\pi(B))$, $\pi : \mathbb{H} \rightarrow \Gamma \backslash \mathbb{H}$ the projection, which follows by applying the Stone-Weierstrass theorem. Note that the uniform closure of the space $\{F_\phi : \phi \in C_c^\infty(\mathbb{H})\}$ is $C_0(\Gamma \backslash \mathbb{H})$, the continuous functions vanishing at infinity.

Option i) On the support of the test function ϕ_0 , we have $v_k \rightarrow -\infty$ uniformly, so that there is some K so that for all $k \geq K$ and all $z \in \text{supp } \phi_0$ we have

$$v_k(z) \leq -2H$$

where $H = \max\{\text{Im } z : z \in \text{supp } \phi_0\}$. But then exponentiating we find

$$|f_k|^2 \leq e^{-2kH}$$

so that for all ϕ supported inside $\text{supp } \phi_0$ (and so that $\text{supp } \phi$ is contained in a single fundamental domain)

$$\int_{\Gamma \backslash \mathbb{H}} \phi(z) |f_k(z)|^2 y^k \frac{dx dy}{y^2} \rightarrow 0$$

contradicting the assumption that $y^k |f_k|^2 dV \xrightarrow{w} cdV$.

Option ii) We assume $\{v_k\}$ has a weakly convergent subsequence, which for notational convenience we continue calling $\{v_k\}$, which then converges to some $v \in \mathcal{SH}$, and moreover $\limsup v_k \leq v$ and these are equal almost everywhere. Then by (9), $v + \log y \leq 0$ almost everywhere.

Since $v_k \xrightarrow{w} v$ but $\int_{\mathbb{H}} v(z) \Delta \phi_0(z) dz \neq \int_{\mathbb{H}} -\log y \Delta \phi_0(z) dz$, we know that $v \neq -\log y$ on a set of positive measure. Since v is upper semi-continuous, there is some $\delta > 0$ so that $v < -\log y - \delta$ on some nonempty open relatively compact set U .

Since $v = \limsup v_k < -\log y - \delta$, by Hartogs' lemma 2.3, there is $K = K(\delta, U)$ so that for all $k \geq K$, $v_k \leq -\log y - \delta/2$ on U . That is to say,

$$\frac{1}{k} \log |f_k|^2 < -\log y - \delta/2$$

or

$$y^k |f_k|^2 \leq e^{-k\delta/2}$$

on U , which as in the first option contradicts $y^k |f_k|^2 dV \xrightarrow{w} cdV$.

4. SKETCH OF THE PROOF OF THEOREM 1

As is well understood by now [16, 21], the equidistribution of masses (1) (with $c = 1$!) follows from the Generalized Riemann Hypothesis (for certain automorphic L -functions), in fact from a “subconvexity estimate” for the central value of certain L -functions. To relate the equidistribution of the masses for Hecke eigenforms to GRH, one has to examine the behavior of the “periods”

$$\int_{SL_2(\mathbb{Z}) \backslash \mathbb{H}} g(z) |f_k(z)|^2 y^k dV(z)$$

where g is fixed and $k \rightarrow \infty$. We need to show that whenever g is orthogonal to the constants then the period vanishes as $k \rightarrow \infty$.

In the case where g is Eisenstein series the period is the (completed) Rankin-Selberg L -function $L^*(1/2 + it, f_k \times f_k)$, the standard convexity estimates give a bound of k^ϵ (for all $\epsilon > 0$) and vanishing as $k \rightarrow \infty$ follows from GRH (see [16, §4]).

To handle the case when g is a cuspidal Hecke-Maass form, one exploits Watson's formula [21] which relates the period with the central value of the triple product L -function:

$$L\left(\frac{1}{2}, g \times f_k \times f_k\right)$$

To get the decay as $k \rightarrow \infty$, it transpires that, again, one has to beat the “convexity bound” on the central value, for which one uses GRH.

In the case of CM-forms³, Sarnak [17] proved that their masses are equidistributed. Thus the reasoning above implies (unconditionally) that the zeros of CM-forms are equidistributed w.r.t. dV as $k \rightarrow \infty$.

APPENDIX A. AN L^∞ -BOUND FOR CUSP FORMS

A holomorphic form of weight k for $SL_2(\mathbb{Z})$ is bounded, since we require that the q -expansion contains no negative powers of q . We will need a bound on the supremum with an explicit dependence on the weight k . For our purposes any bound on $y^{k/2}|f(z)|$ which is sub-exponential in k for fixed z suffices. Below we derive a bound of size $k^{1/2}$; since the dimension of the space of cusp forms grows linearly in k , this bound is optimal (for arbitrary forms) as far as the k -dependence. The proof is adapted from [3, Lemma A.1] who treat the case of Maass forms, though unlike [3] we do not need to control the z -aspect.

A.1. The incomplete Gamma function. We first recall some properties of the incomplete Gamma function $\Gamma(a, x)$ defined for $x > 0$ by

$$\Gamma(a, x) := \int_x^\infty e^{-t} t^a \frac{dt}{t}$$

From the definition it is clear that $\Gamma(a, x)$ is decreasing in x . For integer k we have

$$(10) \quad \Gamma(k-1, x) = (k-2)! e^{-x} \sum_{m=0}^{k-2} \frac{x^m}{m!}$$

What is not completely straightforward is the asymptotic behavior of $\Gamma(a, x)$ when a, x tend to infinity. We will need the asymptotic:

$$(11) \quad \Gamma(k-1, k) \sim \frac{1}{2} \Gamma(k-1), \quad k \rightarrow \infty$$

³These only exist for certain congruence subgroups of the modular group

By (10), the relation (11) is equivalent to

$$(12) \quad e^{-k} \sum_{m=0}^{k-2} \frac{k^m}{m!} \sim \frac{1}{2}$$

This is close to a conjecture of Ramanujan [11, 12, 19, 20] that

$$e^{-k} \left(\sum_{m=0}^{k-1} \frac{k^m}{m!} + \theta \frac{k^k}{k!} \right) = \frac{1}{2}$$

for some $\frac{1}{3} \leq \theta = \theta(k) \leq \frac{1}{2}$; he showed [12] that $\theta(\infty) = 1/3$. We need the weaker asymptotic (12) or equivalently (by Stirling's formula), that

$$(13) \quad e^{-k} \sum_{m=0}^k \frac{k^m}{m!} \sim \frac{1}{2}$$

Here is a quick “proof” using the central limit theorem (this kind of argument goes back to Mark Kac [4]): Let $X_1, \dots, X_k$ be independent Poisson variables with parameter 1 (so having mean and variance 1). Let $S_k = X_1 + \dots + X_k$ be their sum. By the Central Limit Theorem, $\frac{S_k - k}{\sqrt{k}}$ is asymptotically normal, in particular as $k \rightarrow \infty$,

$$\text{Prob}\left(\frac{S_k - k}{\sqrt{k}} \leq 0\right) \rightarrow \frac{1}{2}$$

that is

$$\text{Prob}(S_k \leq k) \rightarrow \frac{1}{2}$$

However since S_k is Poisson with parameter k , we have

$$\text{Prob}(S_k \leq k) = e^{-k} \sum_{m=0}^k \frac{k^m}{m!}$$

which gives (13).

A.2. The supremum of cusp forms.

Proposition 3. *Let f be a cusp form of weight k for $SL_2(\mathbb{Z})$. Then uniformly for z in compact subsets of $\mathbb{H}$,*

$$\frac{y^k |f(z)|^2}{\langle f, f \rangle} \ll k$$

where $\langle f, f \rangle$ is the Peterson inner product.

Proof. Consider the integral of $y^k |f(z)|^2$ over the “Siegel set”

$$\mathfrak{S} := \{z = x + iy : 0 \leq x \leq 1, \quad y > 1/4\pi\}$$

This integral converges since f is exponentially decreasing in the cusp. Since the Siegel set $\mathfrak{S}$ is contained in a fixed number of translates of the standard fundamental domain $\mathcal{F} = \{z = x + iy : |z| \geq 1, |x| \leq 1/2\}$ we have

$$\int_{\mathfrak{S}} |f(z)|^2 y^k \frac{dx dy}{y^2} \ll \langle f, f \rangle$$

On the other hand, using the Fourier expansion $f(z) = \sum_{n \geq 1} a(n) e^{2\pi i n z}$ and Parseval, we find

$$\begin{aligned} \int_{\mathfrak{S}} |f(z)|^2 y^k \frac{dx dy}{y^2} &= \sum_{n \geq 1} |a(n)|^2 \int_{1/4\pi}^{\infty} e^{-4\pi n y} y^{k-2} dy \\ &= \sum_{n \geq 1} \frac{|a(n)|^2}{(4\pi n)^{k-1}} \Gamma(k-1, n) \end{aligned}$$

Thus we find that

$$(14) \quad \sum_{n \geq 1} \frac{|a(n)|^2}{(4\pi n)^{k-1}} \Gamma(k-1, n) \ll \langle f, f \rangle$$

To estimate $y^k |f(z)|^2$, use the Fourier expansion and Cauchy-Schwartz:

$$\begin{aligned} y^k |f(z)|^2 &= y^k \left| \sum_{n \geq 1} a(n) e^{-2\pi n y} e(n x) \right|^2 \\ &\leq \sum_{n \geq 1} \frac{|a(n)|^2}{(4\pi n)^{k-1}} \Gamma(k-1, n) \times y^k \sum_{n \geq 1} e^{-4\pi n y} \frac{(4\pi n)^{k-1}}{\Gamma(k-1, n)} \end{aligned}$$

For the first sum, use (14) to bound it by $\langle f, f \rangle$. Thus we find

$$\frac{y^k |f(z)|^2}{\langle f, f \rangle} \ll y^k \sum_{n \geq 1} e^{-4\pi n y} \frac{(4\pi n)^{k-1}}{\Gamma(k-1, n)}$$

To bound the sum, we split it up into “small” and “large” n ’s. For “small” n , that is those satisfying $n \leq k$, we can give an upper bound as follows: First use $\Gamma(k-1, n) \geq \Gamma(k-1, k)$, for $n \leq k$. This gives

$$\begin{aligned} y^k \sum_{n \leq k} e^{-4\pi n y} \frac{(4\pi n)^{k-1}}{\Gamma(k-1, n)} &\leq \frac{y^k}{\Gamma(k-1, k)} \sum_{n \leq k} e^{-4\pi n y} (4\pi n)^{k-1} \\ &\leq \frac{y}{\Gamma(k-1, k)} \sum_{n=1}^{\infty} e^{-4\pi n y} (4\pi n y)^{k-1} \end{aligned}$$

We have⁴

$$\begin{aligned} \sum_{n=1}^{\infty} e^{-4\pi ny} (4\pi ny)^{k-1} &\leq \int_0^{\infty} e^{-4\pi xy} (4\pi xy)^{k-1} dx + 2e^{-(k-1)}(k-1)^{k-1} \\ &\ll \Gamma(k) \left(\frac{1}{y} + \frac{1}{\sqrt{k}} \right) \end{aligned}$$

Thus we find

$$y^k \sum_{n \leq k} e^{-4\pi ny} \frac{(4\pi n)^{k-1}}{\Gamma(k-1, n)} \ll \frac{y\Gamma(k)}{\Gamma(k-1, k)} \left(\frac{1}{y} + \frac{1}{\sqrt{k}} \right)$$

which by (11) is

$$\ll k \left(1 + \frac{y}{\sqrt{k}} \right)$$

For the sum over the remaining n 's, use (10) to get

$$\Gamma(k-1, n) \geq e^{-n} n^{k-2}$$

which gives

$$\begin{aligned} y^k \sum_{n > k} e^{-4\pi ny} \frac{(4\pi n)^{k-1}}{\Gamma(k-1, n)} &\leq y^k \sum_{n > k} e^{-4\pi ny} \frac{(4\pi n)^{k-1}}{e^{-n} n^{k-2}} \\ &\leq (4\pi y)^k \sum_{n > k} e^{-n(4\pi y - 1)} n \\ &\ll (4\pi y)^k \frac{(k+1)e^{-(k+1)(4\pi y - 1)}}{(1 - e^{-(4\pi y - 1)})^2} \\ &\ll k e^{-4\pi y} \left(\frac{4\pi y e}{e^{4\pi y}} \right)^k \end{aligned}$$

which, for fixed $y \geq \sqrt{3}/2$, is negligible. $\square$

REFERENCES

- [1] J.A. Getz, *A generalization of a theorem of Rankin and Swinnerton-Dyer on zeros of modular forms*. Proc. Amer. Math. Soc. **132** (2004), no. 8, 2221–2231.
- [2] L. Hörmander *The analysis of linear partial differential operators. I. Distribution theory and Fourier analysis*, Second edition. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 256. Springer-Verlag, Berlin, 1990.
- [3] H. Iwaniec and P. Sarnak *L^∞ norms of eigenfunctions of arithmetic surfaces*. Ann. of Math. (2) **141** (1995), no. 2, 301–320.

⁴If $f(t)$ is increasing for $t < t_0$ and decreasing for $t > t_0$ then $\sum_{n=1}^{\infty} f(n) \leq \int_0^{\infty} f(t) dt + 2f(t_0)$

- [4] M. Kac, *Note on the partial sums of the exponential series*. Univ. Nac. Tucumán. Revista A. 3, (1942). 151–153. reprinted in: *Mark Kac: probability, number theory, and statistical physics*. Selected papers. Edited by K. Baclawski and M. D. Donsker. Mathematicians of Our Time, 14. MIT Press, Cambridge, Mass.-London, 1979.
- [5] E. Lindenstrauss, *Invariant measures and arithmetic quantum unique ergodicity*, Annals of Math., to appear
- [6] W. Luo and P. Sarnak *Quantum ergodicity of eigenfunctions on $\mathrm{PSL}_2(\mathbf{Z}) \backslash \mathbf{H}^2$* , Inst. Hautes Études Sci. Publ. Math. No. **81**, (1995), 207–237.
- [7] D. Mumford *Curves and their Jacobians*, The University of Michigan Press, Ann Arbor, Mich., 1975
- [8] A. Neeman *The distribution of Weierstrass points on a compact Riemann surface*, Ann. of Math. (2) **120** (1984), no. 2, 317–328.
- [9] S. Nonnenmacher and A. Voros *Chaotic eigenfunctions in phase space*, J. Statist. Phys. **92** (1998), no. 3–4, 431–518.
- [10] B. Olsen *On higher order Weierstrass points*, Ann. of Math. (2) **95** (1972), 357–364.
- [11] S. Ramanujan, *Question 294*, J. Indian Math. Soc. **3** (1911) 128;
- [12] S. Ramanujan, *On Question 294*, J. Indian Math. Soc. **4** (1912) 151–152.
- [13] F. K. C. Rankin and H. P. F. Swinnerton-Dyer *On the zeros of Eisenstein series*, Bull. London Math. Soc. **2** (1970), 169–170.
- [14] R. A. Rankin *The zeros of certain Poincaré series*, Compositio Math. **46** (1982), no. 3, 255–272.
- [15] T. Ransford *Potential theory in the complex plane*, London Mathematical Society Student Texts, 28. Cambridge University Press, Cambridge, 1995.
- [16] P. Sarnak *Arithmetic quantum chaos*. The Schur lectures (1992) (Tel Aviv), 183–236, Israel Math. Conf. Proc., 8, Bar-Ilan Univ., Ramat Gan, 1995.
- [17] P. Sarnak *Estimates for Rankin-Selberg L -functions and quantum unique ergodicity*. J. Funct. Anal. **184** (2001), no. 2, 419–453.
- [18] B. Shiffman and S. Zelditch *Distribution of zeros of random and quantum chaotic sections of positive line bundles*, Comm. Math. Phys. **200** (1999), no. 3, 661–683.
- [19] G. Szegő, *Über einige von S. Ramanujan gestellte Aufgaben*, J. London Math. Soc. 3 (1928) 225–232.
- [20] G. N. Watson, *Theorems stated by Ramanujan (V): approximations connected with e^x* , Proc. London Math. Soc. (2) **29** (1929) 293–308.
- [21] T. Watson, *Rankin triple products and quantum chaos*, Annals of Math, to appear.

RAYMOND AND BEVERLY SACKLER SCHOOL OF MATHEMATICAL SCIENCES,
TEL AVIV UNIVERSITY, TEL AVIV 69978, ISRAEL (rudnick@math.tau.ac.il)